


NASA TECHNICAL
TRANSLATION



NASA TT F-255


NASA TT F-255



EXPANSION OF POLYNOMIALS IN BESSEL OPERATORS

by S. I. Osipov

From *Zhurnal Vychislitel'noy Matematiki i
Matematicheskoy Fiziki*, Vol. 4, 1964.



NATIONAL AERONAUTICS AND SPACE ADMINISTRATION • WASHINGTON, D. C. • SEPTEMBER 1965



EXPANSION OF POLYNOMIALS IN BESSEL OPERATORS

By S. I. Osipov

Translation of "Razlozheniye mnogochlena ot operatorov Besselya."
Zhurnal Vychislitel'noy Matematiki i Matematicheskoy Fiziki,
Vol. 4, pp. 149-155, 1964.

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

For sale by the Clearinghouse for Federal Scientific and Technical Information
Springfield, Virginia 22151 - Price \$1.00

EXPANSION OF POLYNOMIALS IN BESSEL OPERATORS

S. I. Osipov

Let $D = d/dt$ be a differential operator, $B_k = ((1/t) (tD))^k a$ /149
Bessel operator (see [1]), $k = 2, 3, \dots$, and let $P_n(\lambda) = \lambda^n +$
 $+ a_1 \lambda^{n-1} + \dots + a_n$ be a polynomial with arbitrary constant coeffi-
cients.

In this paper we will expand the operator $P_n(B_k)$ in powers of
 D^* and we apply this expansion to investigation of ordinary linear
differential equations.

Let G_n be a set of functions of t that have n -th order deriva- /150
tives, $n = 1, 2, \dots$, and assume that

$$\begin{aligned} \Phi(n, k, D) &= \sum_{v=1}^{nk} Q(n, k, v, t) D^v + a_n, \\ Q(n, k, v, t) &= \sum_{p=E((v+k-1)/k)}^{\min(v, n)} C(k, v, p) a_{n-p} t^{v-p}, \end{aligned}$$

*The problem of expanding operators coinciding with Bessel operators or operators with similar structure has been discussed in a number of papers by other authors. See, for example, 1) Issledovaniya po integro-differentsial'nym uravneniyam v Kirgizii (Investigations of integro-differential equations in Kirgiz), Vol. II, Frunze, 1962, pp. 300, 310-311; 2) Materialy sed'moy nauchnoy konferentsii kafedry vysshey matematiki Frunzenskogo politekhnicheskogo instituta (Proceedings of the Seventh Scientific Conference of the Departments of Higher Mathematics of Frunze Polytechnic Institute, 1963, pp. 87-90; 3) J. Math. Anal. and Appl., Vol. 6, No. 3, 1963, pp. 395-397; 4) J. Riordan. Introduction to Combinatorial Analysis [Russian translation], For. Lit. Publ. House, Moscow, p. 57, problem 18.

$$\left. \begin{aligned}
C(k, v, p) &= \frac{1}{[(v-p)!]^2} \sum_{m=\max(v-p, p)}^{p(k-1)} \frac{m! S(k, p, m)}{(m+p-v)!}, \\
S(k, p, m) &= (-1)^{pk} \sum_{r=p}^m \frac{(-1)^{r-p} m! (r!)^{k-2}}{[(r-p)!]^{k-1} (m-r)!}, \\
E\left(\frac{v+k-1}{k}\right) &= \text{integral part of } \frac{v+k-1}{k}, \quad a_0 = 1.
\end{aligned} \right\} (1')$$

$$(1)$$

We have the following

Theorem. On the set G_{nk} the operators $P_n(B_k)$ and $\Phi(n, k, D)$ are equivalent, i.e., for any function $g(t) \in G_{nk}$,

$$P_n(B_k) g(t) = \Phi(n, k, D) g(t). \quad (2)$$

Proof. It is not difficult to use induction to prove that the following equation is true for the set G_{pk} :

$$B_k^p = \sum_{v=p}^{pk} A(k, v, p) t^{v-p} D^v, \quad (3)$$

where $A(k, v, p)$ are coefficients that are independent of \underline{t} .

In order to determine these coefficients, we introduce the following polynomial of degree $p(k-1)$:

$$W_{p,k}(t) = e^{\underline{t} B_k^p} e^{-t}. \quad (4)$$

Let

$$L_m(t) = \sum_{r=0}^m \frac{(-1)^r m!}{(r!)^2 (m-r)!} t^r$$

be an m -th order Laguerre polynomial (see [2], p. 110).
It is clear that

$$\int_0^{\infty} e^{-t} L_m(t) W_{p,k}(t) dt = 0 \quad (5)$$

when $m > p(k-1)$.
It is not difficult to obtain

$$\int_0^{\infty} e^{-t} L_m(t) W_{p,k}(t) dt = (-1)^{pk} \int_0^{\infty} e^{-t} B_{k,m}^p L_m(t) dt$$

by integrating by parts and, consequently,

$$\int_0^{\infty} e^{-t} L_m(t) W_{p,k}(t) dt = \begin{cases} 0, & \text{if } m < p, \\ (-1)^p S(k, p, m), & \text{if } p \leq m. \end{cases} \quad (6)$$

By virtue of (5) and (6),

$$W_{p,k}(t) = \sum_{m=p}^{p(k-1)} (-1)^p S(k, p, m) L_m(t),$$

i.e.,

$$W_{p,k}(t) = \sum_{m=p}^{p(k-1)} \sum_{r=0}^m \frac{(-1)^{r+p} S(k, p, m) m!}{(r!)^2 (m-r)!} t^r. \quad (7)$$

/151

The function

$$\delta(m, p) = \begin{cases} 0, & \text{if } m < p, \\ 1, & \text{if } m \geq p, \end{cases} \quad (8)$$

can be used to represent Eq. (7) in the form

$$W_{p,k}(t) = \sum_{m=0}^{p(k-1)} \sum_{r=0}^m \frac{(-1)^{r+p} \delta(m, p) S(k, p, m) m!}{(r!)^2 (m-r)!} t^r,$$

and a change in the order of summation yields

$$W_{p,k}(t) = \sum_{r=0}^{p(k-1)} \sum_{m=r}^{p(k-1)} \frac{(-1)^{r+p} \delta(m, p) S(k, p, m) m!}{(r!)^2 (m-r)!} t^r.$$

The substitution $r = v - p$ leads to the equation

$$W_{p,k}(t) = \sum_{v=p}^{pk} \sum_{m=v-p}^{p(k-1)} \frac{(-1)^v \delta(m, p) S(k, p, m) m!}{[(v-p)!]^2 (m+p-v)!} t^{v-p}. \quad (9)$$

It follows from (3), (4), and (9) that

$$A(k, v, p) = \sum_{m=v-p}^{p(k-1)} \frac{\delta(m, p) S(k, p, m) m!}{[(v-p)!]^2 (m+p-v)!},$$

i.e.,

$$A(k, v, p) = C(k, v, p). \quad (10)$$

In order to complete the proof of the theorem, note that by virtue of (3) and (10), we have

$$P_n(B_k) - a_n = \sum_{p=1}^n \sum_{v=p}^{pk} C(k, v, p) a_{n-p} t^{v-p} D^v.$$

The function (8) can be used to represent this last expression in the form

$$P_n(B_k) - a_n = \sum_{p=1}^{nk} \sum_{v=p}^{nk} \delta(n, p) \delta(pk, v) C(k, v, p) a_{n-p} t^{v-p} D^v,$$

and, changing the order of summation, we obtain

$$P_n(B_k) - a_n = \sum_{v=1}^{nk} \sum_{p=1}^v \delta(n, p) \delta(pk, v) C(k, v, p) a_{n-p} t^{v-p} D^v.$$

It only remains to note that

$$\sum_{p=1}^v \delta(n, p) \delta(pk, v) C(k, v, p) a_{n-p} t^{v-p} = Q(n, k, v, t),$$

which completes the proof.

We shall consider the special case $k = 2$ in more detail. Here we are dealing with the operator $(d/dt) t (d/dt)$, and the operator $\Phi(n, 2, D)$ can be used to expand a polynomial in this operator.

Noting that

$$C(2, v, p) = \frac{p! S(2, p, p)}{[(v-p)!]^2 (2p-v)!}, \quad S(2, p, p) = p!,$$

we find that

/152

$$C(2, v, p) = \frac{(p!)^2}{[(v-p)!]^2 (2p-v)!}, \quad (11)$$

and

$$\Phi(n, 2, D) = \sum_{v=1}^{2n} Q(n, 2, v, t) D^v + a_n,$$

so that

$$Q(n, 2, v, t) = \sum_{p=E((v+1)/2)}^{\min(v,n)} \frac{(p!)^2 a_{n-p}}{[(v-p)!]^2 (2p-v)!} t^{v-p}.$$

This result also follows from the theorem discussed in [3]* when we apply it to the operator $(d/dt) t (d/dt)$.

Consider the coefficients $C(k, v, p)$, defined by formulas (1'). When $k = 2$ these coefficients, as (11) shows, have a simple structure, and it is not difficult to evaluate them. In the general case ($k = 3, 4, \dots$) formulas (1') are rather complex. As a result, it may prove useful to use the following algorithm for evaluation of the coefficients $C(k, v, p)$:

$$C(k, p, p) = (p!)^{k-1}, \quad C(k, p+1, p) = [(p+1)!]^{k-1} - C(k, p, p),$$

and, more generally, for $v = p+1, p+2, \dots, pk$,

*It should be borne in mind that the notation used in this paper is different from that used in [3]. The author of [3] used B_α to denote the operator $t^{-\alpha} D t^{1+\alpha} D$.

$$C(k, v, p) = \frac{(v!)^{k-1}}{[(v-p)!]^k} - \frac{1}{1!} C(k, v-1, p) - \frac{1}{2!} C(k, v-2, p) - \dots$$

(12)

$$\dots - \frac{1}{(v-p)!} C(k, p, p)$$

for $k = 2, 3, \dots$ and $p = 1, 2, \dots$.

We shall now prove (12). By (10) and (3), the coefficients $c(k, v, p)$ are numbers such that for any function $g(t) \in G_{pk}$

$$B_k^p g(t) = \sum_{v=p}^{pk} C(k, v, p) t^{v-p} D^v g(t). \quad (13)$$

We take the function $I_{0,0,\dots,0}^{(k-1)}(k\sqrt[k]{t})$ from [1] as $g(t)$. Since it satisfies the differential equation

$$\frac{1}{t} \left(t \frac{d}{dt} \right)^k y = y,$$

we have

$$B_k^p I_{0,0,\dots,0}^{(k-1)}(k\sqrt[k]{t}) = I_{0,0,\dots,0}^{(k-1)}(k\sqrt[k]{t}), \quad (14)$$

so that the left side of this equation can be treated as the result of p -fold application of the operator $(d/dt)(t(d/dt))^{k-1}$ to the function $I_{0,0,\dots,0}^{(k-1)}(k\sqrt[k]{t})$. Noting that

$$I_{0,0,\dots,0}^{(k-1)} (k \sqrt[k]{t}) = \sum_{r=0}^{\infty} \frac{t^r}{(r!)^k},$$

we can derive the following expression from (13) and (14):

$$\sum_{v=p}^{pk} C(k, v, p) t^{v-p} D^v \sum_{r=0}^{\infty} \frac{t^r}{(r!)^k} = \sum_{r=0}^{\infty} \frac{t^r}{(r!)^k}. \quad (15)$$

But

/153

$$\begin{aligned} \sum_{v=p}^{pk} C(k, v, p) t^{v-p} D^v \sum_{r=0}^{\infty} \frac{t^r}{(r!)^k} &= \sum_{r=0}^{\infty} \left\{ \sum_{v=p}^{pk} C(k, v, p) t^{v-p} D^v \right\} \frac{t^r}{(r!)^k} = \\ &= \sum_{r=p}^{\infty} \left\{ \sum_{v=p}^{\min(r, pk)} C(k, v, p) \frac{1}{(r-v)!} \right\} \frac{t^{r-p}}{(r!)^{k-1}} = \\ &= \sum_{r=0}^{\infty} \left\{ \sum_{v=p}^{\min(r+p, pk)} C(k, v, p) \frac{1}{(r+p-v)!} \right\} \frac{t^r}{[(r+p)!]^{k-1}}, \end{aligned}$$

and comparison of the coefficients of the first $p(k-1) + 1$ powers of t in the right and left sides of (15) gives us

$$\sum_{\mu=p}^{r+p} C(k, \mu, p) \frac{1}{(r+p-\mu)!} = \frac{[(r+p)!]^{k-1}}{(r!)^k} \quad (r = 0, 1, 2, \dots, pk-p).$$

Substitution of $r+p$ by v yields

$$\sum_{\mu=p}^v C(k, \mu, p) \frac{1}{(v - \mu)!} = \frac{(v!)^{k-1}}{[(v - p)!]^k} \quad (v = p, p + 1, \dots, pk),$$

and, consequently, formulas (12) are proved.

The operational calculus of Bessel operators can be used (under certain conditions) to integrate differential equations whose left side is of the form $P_n(B_k)x(t)$. But, both in the theory of ordinary

differential equations and in practice, the left sides of differential equations are usually given in the form of a linear differential expression, i.e., in the form

$$q_0(t) \frac{d^m}{dt^m} x(t) + q_1(t) \frac{d^{m-1}}{dt^{m-1}} x(t) + \dots + q_{m-1}(t) \frac{d}{dt} x(t) + q_m(t) x(t), \quad (16)$$

where $q_0(t), \dots, q_m(t)$ are coefficients and \underline{m} is the order of the differential equation.

In connection with this, it is interesting to investigate the problem of which linear differential equations (16) are representable in terms of $P_n(B_k)x(t)$ and to give a method for reducing (16) to the form $P_n(B_k)x(t)$ when this is possible.

It is a direct consequence of the expansion theorem for $P_n(B_k)$ that representability of (16) in the form $P_n(B_k)x(t)$ requires that $q_0(t), q_1(t), \dots, q_{m-1}(t)$ be polynomials and that $q_m(t)$ be constant.

Let $q_0(t), \dots, q_{m-1}(t)$ be polynomials in \underline{t} and let $q_m(t)$ be a constant, $m \geq 2$ (the cases $m = 0$ and $m = 1$ are trivial). Using the expansion of $P_n(B_k)$ and considering that the number of pairs of integers (n, k) satisfying the conditions $k_n = m$, $k \geq 2$, $n \geq 1$ for any fixed number \underline{m} is finite, we can always establish whether or not (16) is representable in the form $P_n(B_k)x(t)$.

In order to do this we first find all of the admissible pairs of values of \underline{n} and \underline{k} for a given \underline{m} . Among these pairs we select a pair with the largest value of \underline{k} (denote it by k_1 , and denote the corresponding value of \underline{n} by n_1) and we calculate $E((v + k_1 - 1)/k_1)$ and $\min(v, n_1)$ for $v = 1, 2, \dots, m$.

Let $\sigma_{H,v}$ and $\sigma_{B,v}$ be the smallest and largest of the exponents contained in $q_{m-v}(t)$. If $q_{m-v}(t) \equiv 0$, we put

$$\sigma_{H,v} = v - \min(v, n_1), \quad \sigma_{B,v} = v - E\left(\frac{v + k_1 - 1}{k_1}\right).$$

We then construct Table 1 until a negative number appears in column IV or VII. If such a number appears, we conclude that (16) is not representable in the form $P_{n_1}(B_{k_1})x(t)$, and we then turn to testing the next admissible pair of values of \underline{n} and \underline{k} .

TABLE 1

I	II	III	IV	V	VI	VII
v	$\min(v, n_1)$	$\sigma_{H,v}$	$\min(v, n_1) +$ $+ \sigma_{H,v} - v$	$E((v+k_1-1)/k_1)$	$\sigma_{B,v}$	$v - \sigma_{B,v} -$ $-E((v+k_1-1)/k_1)$
1						
.						
.						
.						
.						
m						

If all of the numbers in columns IV and VII are nonnegative, we write the polynomials $q_{m-v}(t)$ in the form

/154

$$\sum_{p=E((v+k_1-1)/k_1)}^{\min(v, n_1)} \beta(v, p) t^{v-p}.$$

TABLE 2

$\begin{array}{c} p \\ \backslash \\ v \end{array}$	1	...	n_1
1	$\beta(1,1)/C(k_1, 1,1)$		
.			
.			
m			
			$\beta(m, n_1)/C(k_1, m, n_1)$

TABLE 3

I	II	III	IV	V	VI	VII
v	$\min(v,1)$	$\sigma_{H,v}$	$\min(v,1)+\sigma_{H,v}$	$E((v+5)/6)$	$\sigma_{B,v}$	$v-\sigma_{B,v}$
1	1	0	0	1	0	$-E((v+5)/6)$
2	1	0	-1			0

TABLE 4

I	II	III	IV	V	VI	VII
v	$\min(v,2)$	$\sigma_{H,v}$	$\min(v,2)+\sigma_{H,v}$	$E((v+2)/3)$	$\sigma_{B,v}$	$v-\sigma_{B,v}$
1	1	0	0	1	0	$-E((v+2)/3)$
2	2	0	0	1	0	0
3	2	1	0	1	1	1
4	2	2	0	2	2	0
5	2	3	0	2	3	0
6	2	4	0	2	4	0

TABLE 5

v \ p	p	
	1	2
1	0	
2	0	1
3	0	1
4		1
5		1
6		1

In this case, $\beta(v, p)$ is uniquely determined by the polynomial $q_{m-v}(t)$. We use formulas (12) to evaluate the coefficients $C(k_1, v, p)$ for $p = 1, 2, \dots, n_1$ and $v = p, p+1, \dots, pk_1$. We place the fractions $\beta(v, p)/C(k_1, v, p)$ at the intersection of the v -th row and the p -th column in Table 2.

When different fractions appear in any column we conclude that (16) is not representable in the form $P_{n_1}(B_{k_1})x(t)$ and turn to testing the next admissible pair of values for \underline{n} and \underline{k} .

If, however, all of the fractions in each column are equal, we conclude that (16) is representable in the form $P_{n_1}(B_{k_1})x(t)$ and that the coefficient a_{n_1-p} , $p = 1, 2, \dots, n_1$, of the polynomial $P_{n_1}(B_{k_1})$ is the number of the p -th column.

Consider, for example, the linear differential expression

$$t^4 \frac{d^6}{dt^6} x(t) + 12t^3 \frac{d^5}{dt^5} x(t) + 38t^2 \frac{d^4}{dt^4} x(t) + 32t \frac{d^3}{dt^3} x(t) + 4 \frac{d^2}{dt^2} x(t). \quad (17)$$

The admissible pairs of values of \underline{n} and \underline{k} are

$$\begin{array}{l} n \ 1 \ 2 \ 3 \\ k \ 6 \ 3 \ 2. \end{array}$$

We first test $n_1 = 1, k_1 = 6$. We construct the first table for this case -- Table 3. At $v = 2$ we obtain a negative number in column IV: -1. As a result, (17) cannot be represented in the form $P_1(B_6)x(t)$.

We now turn to testing the pair $n_2 = 2, k_2 = 3$.

The first table for this case -- Table 4 -- yields no negative entries and we continue the investigation. We have

/155

$$q_5(t) = 0 = \sum_{p=1}^1 \beta(1, p) t^{1-p}, \quad \beta(1,1) = 0;$$

$$q_4(t) = 4 = \sum_{p=1}^2 \beta(2, p) t^{2-p}, \quad \beta(2,1) = 0, \quad \beta(2,2) = 4;$$

$$q_3(t) = 32t = \sum_{p=1}^2 \beta(3, p) t^{3-p}, \quad \beta(3,1) = 0, \quad \beta(3,2) = 32;$$

$$q_2(t) = 38t^2 = \sum_{p=2}^2 \beta(4, p) t^{4-p}, \quad \beta(4,2) = 38;$$

$$q_1(t) = 12t^3 = \sum_{p=2}^2 \beta(5, p) t^{5-p}, \quad \beta(5,2) = 12;$$

$$q_0(t) = t^4 = \sum_{p=2}^2 \beta(6, p) t^{6-p}, \quad \beta(6,2) = 1.$$

Computation of the coefficients $C(3, v, p)$ for $p = 1, 2$, and $v = p, \dots, 3p$ yields

$$c(3, 1, 1) = 1, \quad c(3, 2, 1) = 3, \quad c(3, 3, 1) = 1, \quad c(3, 2, 2) = 4, \\ c(3, 3, 2) = 32, \quad c(3, 4, 2) = 38, \quad c(3, 5, 2) = 12, \quad c(3, 6, 2) = 1,$$

and we obtain Table 5, which shows that (17) is representable in the form $(B_3)^2 x(t)$.

Received: 29 June 1963

BIBLIOGRAPHY

1. Ditkin, V. A. & Prudnikov, A. P. Operatsionnoye ischisleniye operatorov Besselya (Operational Calculus of Bessel Operators), Zhurnal vychislitel'noy matematiki i matematicheskoy fiziki (Journal of Computer Mathematics and Mathematical Physics), Vol. 2, No. 6, 997-1018, 1962.
2. Segë, G. Ortogonal'nyye mnogochleny (Orthogonal Polynomials), Fizmatgiz, Moscow, 1962.
3. Osipov, S. I. O razlozhenii mnogochlena ot operatora B_α (On Expanding Polynomials in B_α Operators), Zh. vychisl. matem. fiz., Vol. 3, No. 1, 191-194, 1963.

FARADAY TRANSLATIONS
15 PARK ROW
NEW YORK, N. Y. 10038

"The aeronautical and space activities of the United States shall be conducted so as to contribute . . . to the expansion of human knowledge of phenomena in the atmosphere and space. The Administration shall provide for the widest practicable and appropriate dissemination of information concerning its activities and the results thereof."

—NATIONAL AERONAUTICS AND SPACE ACT OF 1958

NASA SCIENTIFIC AND TECHNICAL PUBLICATIONS

TECHNICAL REPORTS: Scientific and technical information considered important, complete, and a lasting contribution to existing knowledge.

TECHNICAL NOTES: Information less broad in scope but nevertheless of importance as a contribution to existing knowledge.

TECHNICAL MEMORANDUMS: Information receiving limited distribution because of preliminary data, security classification, or other reasons.

CONTRACTOR REPORTS: Technical information generated in connection with a NASA contract or grant and released under NASA auspices.

TECHNICAL TRANSLATIONS: Information published in a foreign language considered to merit NASA distribution in English.

TECHNICAL REPRINTS: Information derived from NASA activities and initially published in the form of journal articles.

SPECIAL PUBLICATIONS: Information derived from or of value to NASA activities but not necessarily reporting the results of individual NASA-programmed scientific efforts. Publications include conference proceedings, monographs, data compilations, handbooks, sourcebooks, and special bibliographies.

Details on the availability of these publications may be obtained from:

SCIENTIFIC AND TECHNICAL INFORMATION DIVISION
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION
Washington, D.C. 20546